



Stochastic climate theory and modeling

Christian L. E. Franzke,^{1,*} Terence J. O’Kane,² Judith Berner,³
Paul D. Williams⁴ and Valerio Lucarini^{1,5}

Stochastic methods are a crucial area in contemporary climate research and are increasingly being used in comprehensive weather and climate prediction models as well as reduced order climate models. Stochastic methods are used as subgrid-scale parameterizations (SSPs) as well as for model error representation, uncertainty quantification, data assimilation, and ensemble prediction. The need to use stochastic approaches in weather and climate models arises because we still cannot resolve all necessary processes and scales in comprehensive numerical weather and climate prediction models. In many practical applications one is mainly interested in the largest and potentially predictable scales and not necessarily in the small and fast scales. For instance, reduced order models can simulate and predict large-scale modes. Statistical mechanics and dynamical systems theory suggest that in reduced order models the impact of unresolved degrees of freedom can be represented by suitable combinations of deterministic and stochastic components and non-Markovian (memory) terms. Stochastic approaches in numerical weather and climate prediction models also lead to the reduction of model biases. Hence, there is a clear need for systematic stochastic approaches in weather and climate modeling. In this review, we present evidence for stochastic effects in laboratory experiments. Then we provide an overview of stochastic climate theory from an applied mathematics perspective. We also survey the current use of stochastic methods in comprehensive weather and climate prediction models and show that stochastic parameterizations have the potential to remedy many of the current biases in these comprehensive models. © 2014 John Wiley & Sons, Ltd.

How to cite this article:

WIREs Clim Change 2015, 6:63–78. doi: 10.1002/wcc.318

INTRODUCTION

The last few decades have seen a considerable increase in computing power which allows the simulation of numerical weather and climate

prediction models with ever higher resolution and the inclusion of ever more physical processes and climate components (e.g., cryosphere and chemistry). Despite this increase in computer power many important physical processes (e.g., tropical convection, gravity wave drag, and clouds) are still not or only partially resolved in these numerical models. Despite the projected exponential increase in computer power these processes will not be explicitly resolved in numerical weather and climate models in the foreseeable future.^{1,2} For instance, Dawson et al.³ have demonstrated using the ECMWF integrated forecast system that extremely high resolutions (T1279, which corresponds to a grid spacing of about 16 km) are required to simulate accurately the observed Northern hemispheric circulation regime structure. This resolution, typical for limited area weather and climate models

*Correspondence to: christian.franzke@uni-hamburg.de

¹Meteorological Institute and Centre for Earth System Research and Sustainability (CEN), University of Hamburg, Hamburg, Germany

²Centre for Australian Weather and Climate Research, CSIRO Marine and Atmospheric Research, Hobart, Australia

³National Center for Atmospheric Research, Boulder, USA

⁴Department of Meteorology, University of Reading, Reading, UK

⁵Department of Mathematics and Statistics, University of Reading, Reading, UK

Conflict of interest: The authors have declared no conflicts of interest for this article.

used for short-term prediction, remains unfeasible for the current generation of high-resolution global climate models due to computational and data storage requirements. Hence, these missing processes need to be parameterized, i.e., they need to be represented in terms of resolved processes and scales.⁴ This representation is important because small-scale (unresolved) features can impact the larger (resolved) scales^{5,6} and lead to error growth, uncertainty, and biases.

At present, these parameterizations are typically deterministic, relating the resolved state of the model to a unique tendency representing the integrated effect of the unresolved processes. These ‘bulk parameterizations’ are based on the notion that the properties of the unresolved subgrid scales are determined by the large scales. However, studies have shown that resolved states are associated with many possible unresolved states.^{7–9} This calls for stochastic methods for numerical weather and climate prediction that potentially allow a proper representation of the uncertainties, a reduction of systematic biases, and improved representation of long-term climate variability. Furthermore, while current deterministic parameterization schemes are inconsistent with the observed power-law scaling of the energy spectrum^{10,11} new statistical dynamical approaches that are underpinned by exact stochastic model representations have emerged that overcome this limitation. The observed power spectrum structure is caused by cascade processes. Recent theoretical studies suggest that these cascade processes can be best represented by a stochastic non-Markovian Ansatz. Non-Markovian terms are necessary to model memory effects due to model reduction.¹² It means that in order to make skillful predictions the model has to take into account also past states and not only the current state (as for a Markov process).

We first review observational evidence of stochasticity in laboratory geophysical fluid experiments (Section 2), then discuss stochastic climate theory in fast–slow systems (Section 3). In Section 4, we present statistical physics approaches and, in Section 5, we review the current state of stochastic–dynamic weather and climate modeling. We close with an outlook and challenges for the future of weather and climate modeling (Section 6).

LABORATORY EVIDENCE OF STOCHASTICITY

Research on the climate system is somewhat hindered by the obvious difficulties of performing reproducible experiments on the atmosphere and ocean in different parameter regimes. For example, an optical physicist studying the nonlinear response of isolated atoms to

intense electromagnetic waves can easily change the incident wavelength.¹³ In contrast, climate scientists cannot (and arguably should not!) change the rotation rate of the planet or the intensity of the incoming solar radiation. To some extent, numerical simulations come to the rescue, by allowing us to perform virtual experiments. However, the grid spacing in climate models is orders of magnitude larger than the smallest energized scales in the atmosphere and ocean, introducing biases.

Fortunately, there is another option available to us. It is possible to exploit dynamical similarity¹⁴ to study analogs of planetary fluid flow in bespoke laboratory experiments. The traditional set-up is the classic rotating annulus, which has been used for decades to study baroclinic instability and other large-scale phenomena.¹⁵ Recent observations of small-scale inertia–gravity waves embedded within a large-scale baroclinic wave^{16–18} have allowed the scale interactions between these two modes to be studied in a laboratory fluid for the first time. The experimental apparatus consists of a two-layer isothermal annulus forced by a differentially rotating lid, which drives a shear across the internal interface and represents the mid-latitude tropospheric wind shear.

The large-scale baroclinic wave in these laboratory experiments exhibits regime behavior, equilibrating at finite amplitude with a zonal wavenumber of typically 1, 2, or 3. These simple wave modes are regarded as prototypes of the more complicated regime behavior in the atmosphere, such as mid-latitude blocking.^{19,20} A notable finding from repeated experiments using this apparatus is that small-scale inertia–gravity waves can induce large-scale regime transitions, despite the separation of wavelengths by an order of magnitude.²¹ An example of this process is illustrated in Figure 1. A wavenumber 2 mode *without* co-existing inertia–gravity waves (upper row) remains a wavenumber 2 mode indefinitely, drifting around the annulus with the zonal-mean flow. In contrast, with the same parameter values, a wavenumber 2 mode *with* co-existing inertia–gravity waves (lower row) is found to have a finite probability of transitioning to a wavenumber 1 mode. The amplitude of the inertia–gravity waves is controlled here without directly affecting the large-scale mode, by slightly varying the interfacial surface tension between the two immiscible fluid layers.

The laboratory transitions discussed above are reminiscent of noise-induced transitions between different equilibrium states in a meta-stable dynamical system.²⁴ To test this interpretation, a quasi-geostrophic numerical model that captures




FIGURE 1 | Regime transitions in a rotating two-layer annulus laboratory experiment, viewed from above. Different colors correspond to different internal interface heights, through the use of a sophisticated visualization technique.²² In the upper row, small-scale inertia–gravity waves are absent and large-scale regime transitions do not occur. In the lower row, small-scale inertia–gravity waves are present locally in the troughs of the large-scale wave and a large-scale regime transition does occur (from the laboratory experiments of Williams et al.^{17,18,21,23}).

the meta-stability of the large-scale flow in the rotating annulus²⁵ was run with and without weak stochastic forcing added to the potential vorticity evolution equation for each fluid layer. The stochastic forcing was an approximate representation of the inertia–gravity waves, which are inherently ageostrophic and are therefore forbidden from the quasi-geostrophic model. Consistent with the laboratory experiments, only when the noise term was activated did the numerical simulations exhibit large-scale wave transitions in the equilibrated flow.²³ In further numerical experiments, the noise was found to be able to influence wavenumber selection during the developing baroclinic instability.

In summary, the above laboratory experiments constitute the first evidence in a real fluid that small-scale waves may trigger large-scale regime transitions. In a numerical model in which the small-scale waves were absent, the transitions were captured through the addition of stochastic noise. Note that

the small-scale waves satisfy the dispersion relation for inertia–gravity waves and are therefore coherent in space and time, and yet apparently they are ‘sensed’ by the large-scale flow as if they were random fluctuations. These results have led to a possible interpretation of sudden stratospheric warmings as noise-induced transitions.²⁶ Furthermore, these laboratory results help to motivate the development of stochastic parameterizations in climate models and a more general development of stochastic climate theory.

STOCHASTIC CLIMATE THEORY

Climate is a multi-scale system in which different physical processes act on different temporal and spatial scales.²⁷ For instance, on the micro-scale are turbulent eddies with time scales of seconds to minutes, on the meso-scale is convection with time scales of hours to days, on the synoptic scale are

mid-latitude weather systems and blocking with time scales from days to weeks, and on the large-scale are Rossby waves and teleconnection patterns with time scales of weeks to seasons. And there is the coupled atmosphere–ocean system with time scales of seasons to decades. The crucial point here is that all these processes acting on widely different temporal and spatial scales interact with each other due to the inherent nonlinearity of the climate system. We have shown an illustrative laboratory example for this in the previous section.

For many practical applications, we are only interested in the processes on a particular scale and not in the detailed evolution of the processes at smaller scales. Often the scales of interest are linked to inherently predictable processes, while the smaller scales processes are unpredictable. For instance, in the above laboratory experiment we are interested in the regime behavior and not in the detailed evolution of the inertia–gravity waves. In numerical simulations, the fastest scales, which are typically also the smallest scales, use up the bulk of computing time, slowing down the computation of the processes of actual interest. In numerical weather and climate prediction, many of the small-scale processes are currently not explicitly resolved and will not be in the foreseeable future. This neglect of these processes can lead to biases in the simulations. Because of that the unresolved processes need to be parameterized as demonstrated in the previous section.

Stochastic climate theory is based on the concept of scale separation in space or time. Hasselmann²⁸ was the first to propose to split the state vector \vec{z} into slow climate components \vec{x} and fast weather fluctuations \vec{y} and then to derive an effective equation for the slow climate variables only. In this equation, the effect of the now unresolved variables is partially represented as a noise term. The physical intuition behind this idea is, e.g., that the aggregated effect of ‘fast’ weather fluctuations drives fluctuations in the ‘slower’ ocean circulation. To first order, such a model can explain the ‘red’ spectrum of oceanic variables.^{29,30} It has to be noted that there is no scale separation in the climate system. This lack of time-scale separation introduces non-Markovian (memory) effects and complicates the derivation of systematic parameterizations.

Rigorous mathematical derivations for this approach have been provided by Gottwald and Melbourne,³¹ Khasminsky,³² Kurtz,³³ Melbourne and Stuart,³⁴ Papanicolaou,³⁵ and Pavliotis and Stuart.³⁶ For accessible reviews, see Givon et al.³⁷ and the text book by Pavliotis and Stuart.³⁶ This approach has been applied to climate models by Majda and coworkers.^{38–47} Climate models have the following

general functional form

$$d\vec{z} = \left[\tilde{F} + \tilde{L}\vec{z} + \tilde{B}(\vec{z}, \vec{z}) \right] dt \quad (1)$$

where \tilde{F} denotes an external forcing, \tilde{L} a linear operator, and \tilde{B} a quadratic nonlinear operator. Equation (1) constitutes the form of the dynamical cores of weather and climate prediction models.

Now splitting the state vector \vec{z} into slow \vec{x} and fast \vec{y} components (which amounts to assuming a time scale separation) and assuming that the nonlinear self-interaction of the fast modes $\tilde{B}(\vec{y}, \vec{y})$ can be represented by a stochastic process^{39–42} leads to a stochastic differential equation (SDE). The stochastic mode reduction approach^{39–42} then predicts the functional form of reduced climate models for the slow variable \vec{x} alone:

$$d\vec{x} = \left[F + L\vec{x} + B(\vec{x}, \vec{x}) + M(\vec{x}, \vec{x}, \vec{x}) \right] dt + \sigma_A d\vec{W}_A + \sigma_M(\vec{x}) d\vec{W}_M \quad (2)$$

Structurally new terms are a deterministic cubic term that acts predominantly as nonlinear damping and both additive and multiplicative (state-dependent) noise terms. The fundamentals of stochastic processes and calculus are explained in Box 1. The multiplicative noise and the cubic-term stem from the nonlinear interaction between the resolved and unresolved modes.³⁹

The above systematic procedure allows also a physical interpretation of the new deterministic and stochastic terms.³⁹ The additive noise stems both from the nonlinear interaction amongst the unresolved modes and the linear interaction between resolved and unresolved modes.³⁹

Multiplicative (or state-dependent) noise is important for deviations from Gaussianity and thus extremes. The intuition behind multiplicative noise is as follows: On a windless day, the fluctuations are very small, whereas on a windy day not only is the mean wind strong but also the fluctuations around this mean are large; thus, the magnitude of the fluctuations is dependent on the state of the system.

The first practical attempts at stochastic climate modeling were made using Linear Inverse Models (LIM)^{52–55} and dynamically based linear models with additive white noise forcing.^{56–60} These approaches linearize the dynamics and then add white noise and damping⁵⁴ in order to make the models numerically stable (i.e., the resulting linear operator should only have negative eigenvalues to ensure stability and realizability of the solutions). While these models have encouraging predictive skill, especially for ENSO, they

BOX 1

STOCHASTIC PROCESSES

In contrast to deterministic processes, stochastic processes have a random component. See the books by Lemons⁴⁸ and Gardiner⁴⁹ for intuitive introductions to stochastic processes. Typically, stochastic processes are driven by white noise. White noise is a serially uncorrelated time series with zero mean and finite variance.⁴⁹

An SDE is a combination of a deterministic differential equation and a stochastic process. In contrast to regular calculus, stochastic calculus is not unique; i.e., different discretizations of its integral representation lead to different results even for the same noise realization. The two most important calculi are Ito and Stratonovich. See details in the study by Gardiner.⁴⁹ The physical difference is that Ito calculus has uncorrelated noise forcing while Stratonovich allows for finite correlations between noise increments. Hence, physical systems have to be typically approximated by Stratonovich SDEs. However, it is mathematically straightforward to switch between the two calculi. So one only needs to make a decision at the beginning which calculus is more appropriate for modeling the system under consideration and can then switch to the mathematically more convenient form.

SDEs describe systems in a path-wise fashion. The Fokker–Planck equation (FPE) describes how the probability distribution evolves over time.⁴⁹ Thus, SDEs and the FPE offer two different ways at looking at the same system. The parameters of SDEs and their corresponding FPE are linked; thus, one can use the FPE to estimate the parameters of the corresponding SDE.^{50,51}

can only produce Gaussian statistics and, thus, are less useful for predictions of high impact weather.

Recently, there are encouraging attempts in fitting nonlinear stochastic models to data. These include multi-level regression,^{61,62} fitting the parameters via the FPE,^{50,51} stochastic averaging,^{63,64} optimal prediction,^{65,66} or Markov Chains.⁶⁷ Most of the previous approaches fitted the parameters of the stochastic models without taking account of physical constraints, e.g., global stability. Many studies linearized the dynamics and then added additional damping to obtain numerically stable models.^{68,69,54,60} Majda et al.⁴⁷ developed the nonlinear normal form of stochastic climate models and also physical constraints for parameter estimation. Recent studies use these physical constraints to successfully

derive physically consistent stochastic climate models.^{70–72}

Most of the above approaches are based on an implicit assumption of time-scale separation. However, the climate system has a spectrum with no clear gaps which would provide the basis of scale separation and the derivation of reduced order models. Such a lack of time-scale separation introduces memory effects into the truncated description. Memory effects mean that the equations become non-Markovian and that also past states need to be used in order to predict the next state. This can be explained by considering the interaction between a large-scale Rossby wave with a smaller scale synoptic wave. At some location, the Rossby wave will favor the development of the synoptic wave. Initially, this synoptic wave grows over some days without affecting the Rossby wave. Once the synoptic wave has reached a sufficient large amplitude, it will start affecting the Rossby wave. Now in a reduced order model only resolving the Rossby wave but not the synoptic wave this interaction cannot be explicitly represented. However, because the Rossby wave initially triggered the synoptic wave which then in turn affects it some days later, this can be modeled with memory terms which takes into account that the Rossby wave has triggered at time t_0 an anomaly which will affect it at some time later t_n .

Recently, Wouters and Lucarini⁷³ have proposed to treat comprehensively the problem of model reduction in multi-scale systems by adapting the Ruelle response theory^{74,75} for studying the effect of the coupling between the fast and slow degrees of freedom of the system. This theory has previously been used in a geophysical context to study the linear and nonlinear response to perturbations,^{76,77} which also allows climate change predictions. This approach is based on the *chaotic hypothesis*⁷⁸ and allows the general derivation of the reduced dynamics of the slow variables able to mimic the effect of the fast variables in terms of matching the changes in the expectation values of the observables of the slow variables. The ensuing parametrization includes a deterministic correction, which is a mean field result and corresponds to linear response, a general correlated noise and a non-Markovian (memory) term. These results generalize Eq. (2). In the limit of infinite time-scale separation, the classical results of the averaging method are recovered. Quite reassuringly, the same parametrizations can be found using a classical Mori-Zwanzig approach,¹² which is based on projecting the full dynamics on the slow variables and general mathematical results provide evidence that deterministic, stochastic, and non-Markovian components should constitute the backbone of parameterizations.^{79,80}

Recent studies show improvements over approaches based on time-scale separation.^{73,79,80}

Recent studies have shown that stochastic approaches are also important for the prediction of extreme events and tipping points.^{81–84} Sura⁸⁴ discusses a stochastic theory of extreme events. He especially focuses on deviations from a Gaussian distribution; i.e., skewness and kurtosis, as first measures of extremes. He shows that multiplicative noise plays a significant role in causing non-Gaussian distributions. Franzke⁸¹ shows that both deterministic nonlinearity and multiplicative noise are important in predicting extreme events.

STATISTICAL PHYSICS APPROACHES TO STOCHASTIC CLIMATE THEORY

Significant progress has been achieved in the development of tractable and accurate statistical dynamical closures for general inhomogeneous turbulent flows that are underpinned by exact stochastic models (see Box 2). For an accessible review, see the text book by Heinz.⁸⁵ The statistical dynamical closure theory, pioneered by Kraichnan,⁸⁶ has been recognized as a natural framework for a systematic approach to modeling turbulent geophysical flows. Closure theories such as the Direct Interaction Approximation (DIA),⁸⁶ for homogeneous turbulence and the Quasi-Diagonal Direct Interaction Approximation (QDIA),⁸⁷ for the interaction of mean flows with inhomogeneous turbulence have exact generalized Langevin model representations.⁸⁸ This means that such closures are realizable; i.e., they have non-negative energy.

The first major application of turbulence closures has been the examination of the predictability of geophysical flows. Early approaches applied homogeneous turbulence models to predicting error growth^{89–91} whereas more recent advances by Frederiksen and O’Kane,⁹² O’Kane and Frederiksen,⁹³ building on the pioneering studies of Epstein⁹⁴ and Pitcher,⁹⁵ have enabled predictability studies of inhomogeneous strongly non-Gaussian flows typical of the mid-latitude atmosphere. Turbulence closures have also been used for SSP of the unresolved scales, e.g., eddies in atmospheric and ocean general circulation models. As it is generally only possible to represent the statistical effects of unresolved eddies while their phase relationships with the resolved scales are lost,⁹⁶ statistical dynamical turbulence closures are sufficient to allow SSPs to be formulated in a completely transparent way.^{87,89,97–101} Insights gained through the development of inhomogeneous turbulence closure

BOX 2

CLOSURE PROBLEM

In order to describe the statistical behavior of a turbulent flow, the underlying nonlinear dynamical equations must be averaged. For simplicity, we consider a generic equation of motion with quadratic nonlinearity for homogeneous turbulence, in which the mean field is zero, and the fluctuating part of the vorticity in Fourier space, $\hat{\zeta}_{\mathbf{k}}$, satisfies the equation:

$$\frac{\partial}{\partial t} \hat{\zeta}_{\mathbf{k}}(t) = K_{\mathbf{k}\mathbf{p}\mathbf{q}} \hat{\zeta}_{-\mathbf{p}}(t) \hat{\zeta}_{-\mathbf{q}}(t). \quad (3)$$

where \mathbf{p} and \mathbf{q} are the other wave numbers describing triad interactions, i.e., $\mathbf{k} = (k_x, k_y)$ where $\delta(\mathbf{k} + \mathbf{p} + \mathbf{q}) = 1$ if $\mathbf{k} + \mathbf{p} + \mathbf{q} = 0$ and 0 otherwise. Here $K_{\mathbf{k}\mathbf{p}\mathbf{q}}$ are the interaction or mode coupling coefficients. The correlation between the eddies can now be represented by an equation for the covariance (cumulant in terms of wavenumbers \mathbf{k} and \mathbf{l}) which is found to depend on the third-order cumulant in Fourier space:

$$\frac{\partial}{\partial t} \langle \hat{\zeta}_{\mathbf{k}}(t) \hat{\zeta}_{-\mathbf{l}}(t') \rangle = K_{\mathbf{k}\mathbf{p}\mathbf{q}} \langle \hat{\zeta}_{-\mathbf{p}}(t) \hat{\zeta}_{-\mathbf{q}}(t) \hat{\zeta}_{-\mathbf{l}}(t') \rangle. \quad (4)$$

Similarly the third-order cumulant depends on the fourth order and so on, such that we see that an infinite hierarchy of moment or cumulant equations is produced. Statistical turbulence theory is principally concerned with the methods by which this moment hierarchy is closed and the subsequent dynamics of the closure equations. The fact that for homogeneous turbulence the covariance matrix is diagonal greatly simplifies the problem. The majority of closure schemes are derived using perturbation expansions of the nonlinear terms in the primitive dynamical equations. The most successful theories use formal renormalization techniques.^{12,85}

theory have motivated the recent development of general stochastic forms for SSPs for geophysical flows.¹⁰²

Statistical Dynamical Closure Theory

The development of renormalized turbulence closures was pioneered by Kraichnan’s Eulerian DIA⁸⁶ for homogeneous turbulence. The DIA, so named because it only takes into account directly interacting

modes, can be readily regularized to include approximations to the indirect interactions^{103,104} that are required to obtain the correct inertial range-scaling laws. Other homogeneous closures such as Herring's self-consistent field theory (SCFT¹⁰⁵) and McComb's local energy transfer theory (LET),¹⁰⁶ were independently developed soon after. The DIA, SCFT, and LET theories have since been shown to form a class of homogeneous closures that differ only in whether and how a fluctuation dissipation theorem (FDT, i.e., the linear response of a system to an infinitesimal perturbation as it relaxes toward equilibrium)^{86,107–109} is applied. As noted earlier, a consequence of the DIA having an exact stochastic model representation is that it is physically realizable, ensuring positive energy spectra. This is in contrast with closures based on the quasi-normal hypothesis that require further modifications in order to ensure realizability; an example of such a closure is the eddy damped quasi-normal Markovian (EDQNM) closure^{89,110,111} developed as a best Markovian fit to the DIA. The EDQNM is dependent on a choice of an eddy-damping parameter that can be tuned to match the phenomenology of the inertial range. This Markovian assumption assumes that the rate at which the memory integral decays is significantly faster than the time scale on which the covariances evolve. The relative success of these turbulence closures has enabled the further study of the statistics of the predictability of homogeneous turbulent flows.^{89,90,112,113}

Frederiksen⁸⁷ formulated a computationally tractable non-Markovian (memory effects) closure, the QDIA, for the interaction of general mean and fluctuating flow components with inhomogeneous turbulence and topography. The QDIA assumes that, prior to renormalization, a perturbative expansion of the covariances are diagonal at zeroth order. In general, very good agreement has been found between the QDIA closure results and the statistics of direct numerical simulation (DNS).^{92,103,104}

The non-Markovian closures discussed above are systems of integro-differential equations with potentially long-time history integrals posing considerable computational challenges; however, various ways to overcome these challenges exist^{92,93,100,104,109,114–116} and have been generalized to allow computationally tractable closure models for inhomogeneous turbulent flow over topography to be developed.^{87,92,104} An alternative derivation of a stochastic model of the Navier–Stokes equations has been put forward by Memin.¹¹⁷ It is based on a decomposition of the velocity fields into a differentiable drift and a stochastic component.

Statistical Dynamical and Stochastic Subgrid Modeling

Many subgrid-scale stress models assume the small scales to be close to isotropic and in equilibrium such that energy production and dissipation are in balance, similar to the eddy viscosity assumption of the Smagorinsky model.¹¹⁸ Using the DIA, Kraichnan⁸⁶ showed that for isotropic turbulence the inertial transfer of energy could be represented as a combination of both an eddy viscous (on average energy drain from retained to subgrid scales) and stochastic back-scatter (positive semi-definite energy input from subgrid to retained scales) terms. The nonlinear transfer terms represented by eddy viscosity and stochastic back-scatter are the subgrid processes associated with the respective eddy-damping and nonlinear noise terms that constitute the right-hand side of the DIA tendency equation for the two-point cumulant $\frac{\partial}{\partial t} \langle \hat{\zeta}_k(t) \hat{\zeta}_{-k}(t') \rangle$. Leith⁸⁹ used the EDQNM closure to calculate an eddy dissipation function that would preserve a stationary k^{-3} kinetic energy spectrum for two-dimensional turbulence. Kraichnan⁹⁸ developed the theory of eddy viscosity in two and three dimensions and was the first to identify the existence of a strong cusp in the spectral eddy viscosity near the cutoff wavenumber representing local interactions between modes below and near the cusp. Rose¹⁰¹ argued for the importance of eddy noise in subgrid modeling.

O'Kane and Frederiksen¹⁰⁰ calculated QDIA-based SSPS considering observed atmospheric flows over global topography and quantifying the relative importance of the subgrid-scale eddy-topographic, eddy-mean field, quadratic mean, and mean field-topographic terms. They also compared the QDIA-based SSPS to heuristic approaches based on maximum entropy, used to improve systematic deficiencies in ocean climate models.¹¹⁹ While closure models may be the natural starting place for developing SSPs, their complexity makes them difficult to formulate and apply to multi-field models such as GCMS, even though successful studies exist.^{102,120}

STOCHASTIC PARAMETRIZATION SCHEMES IN COMPREHENSIVE MODELS

Climate and weather predictions are only feasible because the governing equations of motion and thermodynamics are known. To solve these equations, we need to resort to numerical simulations that discretize the continuous equations onto a finite grid and parameterize all processes that cannot be explicitly




FIGURE 2 | Mean systematic error of 500 hPa geopotential height fields (shading) for extended boreal winters (December–March) of the period 1962–2005. Errors are defined with regard to the observed mean field (contours), consisting of a combination of ERA-40 (1962–2001), and operational ECMWF analyses (2002–2005). (a) Systematic error in a numerical simulation with the ECMWF model IFS, version CY32R1, run at a horizontal resolution of $T_{1,95}$ (about 210 km) and 91 vertical levels. (b) Systematic error in a simulation with a stochastic kinetic-energy backscatter scheme. Significant differences at the 95% confidence level based on a Student's t -test are hatched (after Berner et al.¹²⁴).

resolved. Such models can be characterized in terms of their dynamical core, which describe the resolved scales of motion, and the physical parameterizations, which provide estimates of the grid-scale effect of processes that cannot be resolved by the dynamical core. This general approach has been hugely successful in that nowadays predictions of weather and climate are made routinely. However, exactly through these predictions it has become apparent that uncertainty estimates produced by current state-of-the-art models still have shortcomings.

One shortcoming is that many physical parameterizations are based on bulk formulae that are based on the assumption that the subgrid-scale state is in equilibrium with the resolved state.¹²¹ Model errors might arise from a misrepresentation of unresolved subgrid-scale processes that can affect not only the variability, but also the mean error of a model.^{122,123} An example in a comprehensive climate model is, e.g., the bias in the 500 hPa geopotential height pattern, which is reduced when the representation of the subgrid state is refined¹²⁴ (Figure 2).

In recent years, methods for the estimation of flow-dependent uncertainty in predictions have become an important topic. Ideally, uncertainties should be estimated within the physical parameterizations and uncertainty representations should be developed alongside the model. Many of these methods are *ad hoc* and added *a posteriori* to an already tuned model. Only first steps to develop

uncertainty estimates from within the parameterizations have been attempted.^{125,126}

The representation of model-error in weather and climate models falls in one of two major categories: multi-model approaches and stochastic parameterizations. In the multi-model approach, each ensemble member consists of an altogether different model. The models can differ in the dynamical core and the physical parameterizations^{127–129} or use the same dynamical core but utilize either different static parameters in their physical parameterizations¹³⁰ or altogether different physics packages.^{131–134} Both approaches have been successful in improving predictions of weather and climate over a range of scales, as well as their uncertainty. Multi-model ensembles provide more reliable seasonal forecasts¹³⁵ and are commonly used for the uncertainty assessment of climate change predictions, e.g., as in the Fifth Assessment Report (AR5) of the Intergovernmental Panel on Climate Change (IPCC).¹³⁶ Stochastic parameterizations are routinely used to improve the reliability of weather forecasts in the short-¹³¹ and medium-range^{10,137,138} as well as for seasonal predictions.^{139–141}

In the stochastic approach, the effect of uncertainties due to the finite truncation of the model is treated as independent realizations of stochastic processes that describe these truncation uncertainties. This treatment goes back to the idea of stochastic-dynamic prediction.^{95,121,142} While the verdict is still open if subgrid-scale fluctuations must

be included explicitly via a stochastic term, or if it is sufficient to include their mean influence by improved deterministic physics parameterizations, one advantage of stochastically perturbed models is that all ensemble members have the same climatology and model bias; while for multi-parameter, multi-parameterization, and multi-model ensembles each ensemble member is *de facto* a different model with its own dynamical attractor. For operational centers, the maintenance of different parameterizations requires additional resources and due to the different biases makes post-processing very difficult.

Stochastic Parameterizations in Numerical Weather Prediction

Due to the chaotic nature of the dynamical equations governing the evolution of weather, forecasts are sensitive to the initial condition limiting the intrinsic predictability of the weather system.^{5,143} Probabilistic forecasts are performed by running ensemble systems, where each member is initialized from a different initial state and much effort has gone into the optimal initialization of such ensemble systems.^{128,144,145} Nevertheless state-of-the-art numerical weather predictions systems continue to produce unreliable and over-confident forecasts.¹⁴⁶ Consequently, the other source of forecast uncertainty—model error—has received increasing attention.^{121,147} Since for chaotic systems model-error and initial condition error will both result in trajectories that will diverge from the truth, it is very difficult to disentangle them.¹⁴⁸

The first stochastic parameterization used in an operational numerical weather prediction model was the stochastically perturbed physics tendency scheme (SPPT), sometimes also referred to as stochastic diabatic tendency or Buizza–Miller–Palmer scheme.¹⁴⁹ SPPT is based on the notion that—especially as the horizontal resolution increases—the equilibrium assumption no longer holds and the subgrid-scale state should be sampled rather than represented by the equilibrium mean. Consequently, SPPT multiplies the accumulated physical tendencies of temperature, wind, and moisture at each grid-point and time step with a random pattern that has spatial and temporal correlations. In other words, SPPT assumes that parameterization uncertainty can be expressed as a multiplicative noise term. Ensemble systems perturbed with the SPPT scheme show increased probabilistic skill mostly due to increased spread in short- and medium-range ensemble forecasts.^{149–152}

A second successful stochastic parameterization scheme, is the so-called stochastic kinetic energy backscatter scheme (SKEBS) whose origin lies in

large-eddy simulation modeling¹⁵³ and has recently been extended to weather and climate scales.^{11,154} The key idea is that energy associated with subgrid processes is injected back onto the grid using a stochastic pattern generator. This method has been successfully used in a number of operational and research forecasts across a range of scales.^{10,131,137,150,155–157} Similar to the SPPT scheme, ensemble systems with SKEBS increase probabilistic skill by increasing spread and decreasing the root-mean-square (RMS) error of the ensemble mean forecast. First results of these schemes at a convection-permitting resolution of around 4 km report also a positive impact on forecast skill, in particular, more reliable precipitation forecasts.^{158,159}

Stochastic Parameterizations in Climate Models

The use of stochastic parameterization in climate models is still in its infancy. Climate prediction uncertainty assessments, e.g., IPCC,¹⁶⁰ are almost exclusively based on multi-models, mostly from different research centers. Part of the problem is that on climate timescales, limited data for verification exist. A second reason is that on longer timescales, bias is a major source of uncertainty and traditional multi-models are very efficient at sampling biases, although such an experiment is poorly designed for an objective and reliable uncertainty assessment.

However, in recent years, first studies have emerged which demonstrate the ability of stochastic parameterizations to reduce longstanding biases and improve climate variability (see Figure 2 for an example). Jin and Neelin¹⁶¹ developed a stochastic convective parameterization that includes a random contribution to the convective available potential energy in the deep convective scheme. They find that adding convective noise results in enhanced eastward propagating, low-wavenumber low-frequency variability. Berner et al.¹²⁴ investigate the impact of SKEBS on systematic model error and report an improvement in the representation of convectively coupled waves leading to a reduction in the tropical precipitation bias. Furthermore, Majda and colleagues developed systematic stochastic multi-cloud parameterizations for organized convection.^{42,162–164} The multi-cloud approach is based on the assumption that organized convection involves three types of clouds and the evolution from one cloud type to another can be described by a transition matrix.

A longstanding systematic error of climate models is the underestimation of the occurrence of Northern Hemispheric blocking. Stochastic parameterizations have been demonstrated to be one way

to increase their frequency,^{139–141,165} although, e.g., increasing horizontal resolution, leads to similar improvements.^{3,124} This suggests that while it might be necessary to include subgrid-scale variability in some form, the details of this representation might not matter. On the other side, Berner et al.¹²⁴ argue that this degeneracy of response to different subgrid-scale forcings warrants a cautionary note: namely that a decrease in systematic error might not necessarily occur for the right dynamical reasons. The opposite holds true, as well: due to the necessary tuning of parameters in the parameterizations of comprehensive climate models, an improvement in the formulation of a physical process might not immediately lead to an improved model performance. A striking example of compensating model errors is described in Palmer and Weisheimer,¹⁶⁶ who report how an inadequate representation of horizontal baroclinic fluxes resulted in a model error that was equal and opposite to the systematic error caused by insufficiently represented vertical orographic gravity wave fluxes. Improvements to wave drag parameterization without increasing resolution unbalanced the compensating model errors, leading to an increase in systematic model bias.

Williams¹⁶⁷ studied the effect of including a stochastic term in the fluxes between the atmospheric and oceanic components in a coupled ocean–atmosphere model. He reports changes to the time-mean climate and increased variability of the El Niño Southern Oscillation, suggesting that the lack of representation of subgrid variability in air–sea fluxes may contribute to some of the biases exhibited by contemporary climate models.

On seasonal timescales where sufficient observational data for a probabilistic verification exist, stochastic parameterizations have been reported to increase predictive skill. For example, ensemble forecasts of the sea surface temperatures over the Niño3.4 region showed increased anomaly correlation, decreased bias, and decreased RMS error in coupled ocean–atmosphere models.^{139–141}

CONCLUSION

We postulate the use of stochastic–dynamical models for uncertainty assessment and model-error representation in comprehensive Earth System Models. This need arises since even state-of-the-art weather and climate models cannot resolve all necessary processes and scales. Here, we reviewed mathematical methods for stochastic climate modeling as well as stochastic SSPs and postulate their use for a more systematic strategy of parameterizing unresolved and unrepresented processes.

In the last decade, a number of studies emerged that demonstrate the potential of this approach, albeit applied in an ad hoc manner and tuned to specific applications. Stochastic parameterizations have been shown to provide more skillful weather forecasts than traditional ensemble prediction methods, at least on timescales where verification data exists. In addition, they have been shown to reduce longstanding climate biases, which play an important role especially for climate and climate change predictions.

Here we argue that rather than pushing out the limit of skillful ensemble predictions by a few days, more attention should be given to the assessment of uncertainty (as already proposed, e.g., Smith¹⁴⁸). Ideally, it should be carried out alongside the physical parameterization and dynamical core development and not added *a posteriori*. The uncertainty should be directly estimated from within the parameterization schemes and not tuned to yield a particular model performance, as is current practice. For example, Sapsis and Majda^{168,169} propose a statistical framework which systematically quantifies uncertainties in a stochastic fashion.

The fact that according to the last two ARs of the IPCC (AR4¹⁷⁰ and AR5¹⁶⁰) the uncertainty in climate predictions and projections has not decreased may be a sign that we might be reaching the limit of climate predictability, which is the result of the intrinsically nonlinear character of the climate system (as first suggested by Lorenz¹⁴³).

Recently Palmer¹⁷¹ argued that due to limited computational and energy power resources, predictable scales should be solved accurately, while the unpredictable scales could be represented inaccurately. This strategy is at the core of the systematic mode reduction reviewed here, but has only recently been considered for comprehensive Earth System Models. Stochastic models focus on the accurate simulation of the large, predictable scales, while only the statistical properties of the small, unpredictable, scales are captured. This has been demonstrated, e.g. by Franzke and Majda,⁴⁰ Kravtsov et al.,⁶² who successfully applied mode reduction strategies to global atmospheric circulation models. They showed that these reduced models consisting of only 10–15 degrees of freedom reproduced many of the important statistics of the numerical circulation models which contained a few hundreds degrees of freedom. Vanden-Eijnden¹⁷² proposed numerical approaches for multi-scale systems where only the largest scales are explicitly computed and the smaller scales are approximated on the fly.

The recent results of Wouters and Lucarini^{73,80} provide a promising path toward a general theory of

parameterizations for weather and climate models, and give theoretical support that parameterization schemes should include deterministic, stochastic and non-Markovian (memory) components. Moreover, Wouters and Lucarini's results suggest that there is common ground in developing parameterizations for weather and climate prediction models. Optimal representations of the reduced dynamics based on Ruelle's response theory and the Mori-Zwanzig formalism coincide, thus, providing equal optimal representations of the long-term statistical properties and the finite-time evolution of the slow variables.

One exciting future research area is the use of stochastic methods for use in data assimilation, which is already an active field of research.^{114,115,159,173–176} Stochastic methods have been shown to increase

the ensemble spread in data assimilation, leading to a better match between observations and model forecasts.^{159,174,177} A cutting-edge frontier is the use of order moments and memory effects in Kalman filter data assimilation schemes.¹¹⁵ Another emerging field is the use of stochastic parameterizations in large climate ensembles, which would allow the comparison of uncertainty estimates based in multi-models to that of stochastically perturbed ones.

Our hope is that basing stochastic–dynamic prediction on sound mathematical and statistical physics concepts will lead to substantial improvements, not only in our ability to simulate weather and climate accurately, but even more importantly to give proper estimates of the uncertainty of these predictions.

ACKNOWLEDGMENTS

CLEF is supported by the German Research Foundation (DFG) through the cluster of Excellence CliSAP, TJO is an Australian Research Council Future Fellow, PDW acknowledges a University Research Fellowship from the Royal Society (UF080256), and VL obtained funding from the European Research Council (NAMASTE).

REFERENCES

- Palmer TN, Williams PD. Introduction: Stochastic physics and climate modelling. *Phil Trans R Soc A* 2008, 366:2421–2427.
- Williams PD. Modelling climate change: the role of unresolved processes. *Phil Trans R Soc A* 2005, 363:2931–2946.
- Dawson A, Palmer TN, Corti S. Simulating regime structures in weather and climate prediction models. *Geophys Res Lett* 2012, 39:L21805.
- Stensrud DJ. *Parameterization Schemes: Keys to Understanding Numerical Weather Prediction Models*. Cambridge: Cambridge University Press; 2007, 459.
- Lorenz EN. Atmospheric predictability as revealed by naturally occurring analogues. *J Atmos Sci* 1969, 26:636–646.
- Tribbia JJ, Baumhefner DP. Scale interactions and atmospheric predictability: an updated perspective. *Mon Weather Rev* 2004, 132:703–713.
- Crommelin D, Vanden-Eijnden E. Subgrid-scale parameterization with conditional Markov chains. *J Atmos Sci* 2008, 65:2661–2675.
- Shutts GJ, Palmer TN. Convective forcing fluctuations in a cloud-resolving model: relevance to the stochastic parameterization problem. *J Climate* 2007, 20:187–202.
- Wilks DS. Effects of stochastic parameterizations in the Lorenz'96 system. *Q J Roy Meteorol Soc* 2005, 131:389–407.
- Berner J, Shutts G, Leutbecher M, Palmer T. A spectral stochastic kinetic energy backscatter scheme and its impact on flow-dependent predictability in the ECMWF ensemble prediction system. *J Atmos Sci* 2009, 66:603–626.
- Shutts GJ. A kinetic energy backscatter algorithm for use in ensemble prediction systems. *Q J Roy Meteorol Soc* 2005, 612:3079–3102.
- Chorin AJ, Hald OH. *Stochastic Tools in Mathematics and Science*, vol. 208. New York: Springer; 2013.
- O'Kane TJ, Scholten RE, Walkiewicz MR, Farrell PM. Nonlinear interactions of multi-level atoms with a near resonant standing wave. *Phys Rev A* 1999, 59:4485–4493.
- Douglas JF, Gasiorek JM, Swaffield JA, Jack LB. *Fluid Mechanics*. 6th ed. Prentice Hall: Harlow; 2011.
- Hide R, Mason PJ, Plumb RA. Thermal convection in a rotating fluid subject to a horizontal temperature gradient: spatial and temporal characteristics of fully developed baroclinic waves. *J Atmos Sci* 1977, 34:930–950.
- Lovegrove AF, Read PL, Richards CJ. Generation of inertia-gravity waves in a baroclinically unstable fluid. *Q J Roy Meteorol Soc* 2000, 126:3233–3254.
- Williams PD, Haine TWN, Read PL. On the generation mechanisms of short-scale unbalanced modes in rotating two-layer flows with vertical shear. *J Fluid Mech* 2005, 528:1–22.

18. Williams PD, Haine TWN, Read PL. Inertia–gravity waves emitted from balanced flow: observations, properties, and consequences. *J Atmos Sci* 2008, 65:3543–3556.
19. Tian Y, Weeks ER, Ide K, Urbach JS, Baroud CN, Ghil M, Swinney HL. Experimental and numerical studies of an eastward jet over topography. *J Fluid Mech* 2001, 438:129–157.
20. Weeks ER, Tian Y, Urbach JS, Ide K, Swinney HL, Ghil M. Transitions between blocked and zonal flows in a rotating annulus with topography. *Science* 1997, 278:1598–1601.
21. Williams PD, Read PL, Haine TWN. Spontaneous generation and impact of inertia–gravity waves in a stratified, two-layer shear flow. *Geophys Res Lett* 2003, 30:2255.
22. Williams PD, Read PL, Haine TWN. A calibrated, non-invasive method for measuring the internal interface height field at high resolution in the rotating, two-layer annulus. *Geophys Astrophys Fluid Dyn* 2004b, 98:453–471.
23. Williams PD, Haine TWN, Read PL. Stochastic resonance in a nonlinear model of a rotating, stratified shear flow, with a simple stochastic inertia-gravity wave parameterization. *Nonlin Proc Geophys* 2004a, 11:127–135.
24. De Swart HE, Grasman J. Effect of stochastic perturbations on a low-order spectral model of the atmospheric circulation. *Tellus A* 1987, 39:10–24.
25. Williams PD, Haine TWN, Read PL, Lewis SR, Yamazaki YH. QUAGMIRE v1.3: a quasi-geostrophic model for investigating rotating fluids experiments. *Geosci Model Dev* 2009, 2:13–32.
26. Birner T, Williams PD. Sudden stratospheric warmings as noise-induced transitions. *J Atmos Sci* 2008, 65:3337–3343.
27. Klein R. Scale-dependent models for atmospheric flows. *Annu Rev Fluid Mech* 2010, 42:249–274.
28. Hasselmann K. Stochastic climate models: part I, theory. *Tellus* 1976, 28:473–484.
29. Frankignoul C, Hasselmann K. Stochastic climate models, part II: application to sea-surface temperature anomalies and thermocline variability. *Tellus* 1977, 29:289–305.
30. Imkeller P, von Storch J-S. *Stochastic Climate Models*. Birkhäuser; 2001, 398.
31. Gottwald G, Melbourne I. Homogenization for deterministic maps and multiplicative noise. *Proc R Soc A* 2013, 469:20130201.
32. Khasminsky RZ. Principle of averaging for parabolic and elliptic differential equations and for Markov processes with small diffusions. *Theory Probab Appl* 1963, 8:1–21.
33. Kurtz TG. A limit theorem for perturbed operator semi-groups with applications to random evolutions. *J Funct Anal* 1973, 12:55–67.
34. Melbourne I, Stuart AM. A note on diffusion limits of chaotic skew-product flows. *Nonlinearity* 2011, 24:1361–1367.
35. Papanicolaou GC. Some probabilistic problems and methods in singular perturbations. *Rocky Mt J Math* 1976, 6:653–674.
36. Pavliotis, GA and AM Stuart, 2008: *Multiscale Methods: Averaging and Homogenization*. Springer, 307.
37. Givon D, Kupferman P, Stuart A. Extracting macroscopic dynamics: model problems and algorithms. *Nonlinearity* 2004, 17:R55.
38. Dolaptchiev SI, Achatz U, Timofeyev I. Stochastic closure for local averages in the finite-difference discretization of the forced Burgers equation. *Theor Comp Fluid Dyn* 2013, 27:297–317.
39. Franzke C, Majda AJ, Vanden-Eijnden E. Low-order stochastic mode reduction for a realistic barotropic model climate. *J Atmos Sci* 2005, 62:1722–1745.
40. Franzke C, Majda AJ. Low-order stochastic mode reduction for a prototype atmospheric GCM. *J Atmos Sci* 2006, 63:457–479.
41. Majda AJ, Timofeyev I, Vanden-Eijnden E. Models for stochastic climate prediction. *Proc Natl Acad Sci USA* 1999, 96:14687–14691.
42. Majda AJ, Timofeyev I, Vanden-Eijnden E. A mathematical framework for stochastic climate models. *Comm Pure Appl Math* 2001, 54:891–974.
43. Majda AJ, Timofeyev I, Vanden-Eijnden E. A priori tests of a stochastic mode reduction strategy. *Phys D* 2002, 170:206–252.
44. Majda AJ, Timofeyev I, Vanden-Eijnden E. Systematic strategies for stochastic mode reduction in climate. *J Atmos Sci* 2003, 60:1705–1722.
45. Majda AJ, Abramov R and Grote M, Information theory and stochastics for multiscale nonlinear systems. CRM Monograph Series, American Mathematical Society, 2005.
46. Majda AJ, Franzke C, Khouider B. An applied mathematics perspective on stochastic modelling for climate. *Phil Trans R Soc A* 2008, 366:2429–2455.
47. Majda AJ, Franzke C, Crommelin D. Normal forms for reduced stochastic climate models. *Proc Natl Acad Sci USA* 2009, 106:3649–3653. doi: 10.1073/pnas.0900173106.
48. Lemons DS. *An Introduction to Stochastic Processes in Physics*. Baltimore: Johns Hopkins University Press; 2002, 115.
49. Gardiner CW. *Handbook of Stochastic Methods*. Berlin: Springer-Verlag; 1985, 442.
50. Berner J. Linking nonlinearity and non-Gaussianity of planetary wave behavior by the Fokker-Planck equation. *J Atmos Sci* 2005, 62:2098–2117.
51. Siegert S, Friedrich R, Peinke J. Analysis of data sets of stochastic systems. *Phys Lett A* 1998, 243:275–280.

52. Penland C, Matrosova L. A balance condition for stochastic numerical models with application to the El Nino-Southern Oscillation. *J Climate* 1994, 7:1352–1372.
53. Penland C, Sardeshmukh PD. The optimal growth of tropical sea surface temperature anomalies. *J Climate* 1995, 8:1999–2014.
54. Whitaker JS, Sardeshmukh PD. A linear theory of extratropical synoptic eddy statistics. *J Atmos Sci* 1998, 55:237–258.
55. Winkler CR, Newman M, Sardeshmukh PD. A linear model of wintertime low-frequency variability, part I: formulation and forecast skill. *J Climate* 2001, 14:4474–4494.
56. DelSole T, Hou AY. Empirical stochastic models for the dominant climate statistics of a General Circulation Model. *J Atmos Sci* 1999, 56:3436–3456.
57. DelSole T. Stochastic models of quasigeostrophic turbulence. *Surv Geophys* 2004, 25:107–149.
58. Farrell BF, Ioannou PJ. Stochastic dynamics of baroclinic waves. *J Atmos Sci* 1993, 50:4044–4057.
59. Farrell BF, Ioannou PJ. Stochastic dynamics of the midlatitude atmospheric jet. *J Atmos Sci* 1995, 52:1642–1656.
60. Zhang Y, Held IM. A linear stochastic model of a GCM's midlatitude storm tracks. *J Atmos Sci* 1999, 56:3416–3435.
61. Kondrashov D, Kravtsov S, Ghil M. Empirical mode reduction in a model of extratropical low-frequency variability. *J Atmos Sci* 2006, 63:1859–1877.
62. Kravtsov S, Kondrashov D, Ghil M. Multilevel regression modeling of nonlinear processes: derivation and applications to climatic variability. *J Climate* 2005, 18:4404–4424.
63. Culina J, Kravtsov S, Monahan AH. Stochastic parameterization schemes for use in realistic climate models. *J Atmos Sci* 2011, 68:284–299.
64. Monahan AH, Culina J. Stochastic averaging of idealized climate models. *J Climate* 2011, 24:3068–3088.
65. Chorin AJ, Hald OH, Kupferman R. Optimal prediction with memory. *Phys D* 2002, 166:239–257.
66. Stinis P. A comparative study of two stochastic mode reduction methods. *Phys D* 2006, 213:197–213.
67. Crommelin D, Vanden-Eijnden E. Reconstruction of diffusions using spectral data from timeseries. *Commun Math Sci* 2006, 4:651–668.
68. Achatz U, Branstator G. A two-layer model with empirical linear corrections and reduced order for studies of internal climate variability. *J Atmos Sci* 1999, 56:3140–3160.
69. Achatz U, Opsteegh JD. Primitive-equation-based low-order models with seasonal cycle, part I: model construction. *J Atmos Sci* 2003, 60:465–477.
70. Harlim J, Mahdi A, Majda AJ. An ensemble Kalman filter for statistical estimation of physics constrained nonlinear regression models. *J Comput Phys* 2013, 257:782–812.
71. Majda AJ, Harlim J. Physics constrained nonlinear regression models for time series. *Nonlinearity* 2013, 26:201–217.
72. Peavoy D, Franzke C and Roberts G: Systematic physics constrained parameter estimation of stochastic differential equations. *Comp Stat Data Anal* arXiv:1312.1881.
73. Wouters J, Lucarini V. Disentangling multi-level systems: averaging, correlations and memory. *J Stat Mech* 2011, 2012:PO3003.
74. Ruelle D. Smooth dynamics and new theoretical ideas from nonequilibrium statistical mechanics. *J Stat Phys* 1999, 95:393–468.
75. Ruelle D. A review of linear response theory for general differentiable dynamical systems. *Nonlinearity* 2009, 22:855.
76. Lucarini V. Evidence for dispersion relations for the nonlinear response of the Lorenz 1963 system. *J Stat Phys* 2009, 134:381–400.
77. Lucarini V, Sarno S. A statistical mechanic approach for the computation of the climatic response to general forcings. *Nonlin Proc Geophys* 2011, 18:7–28.
78. Gallavotti G, Cohen EGD. Dynamical ensembles in stationary states. *J Stat Phys* 1995, 80:931–970.
79. Chekroun M, Liu H, Wang S, On stochastic parameterizing manifolds: pullback characterization and non-Markovian reduced equations. *Mem Am Math Soc* 2013, arxiv:1310.3896.
80. Wouters J, Lucarini V. Multi-level dynamical systems: connecting the Ruelle response theory and the Mori-Zwanzig approach. *J Stat Phys* 2013, 151:850–860.
81. Franzke C. Predictability of extreme events in a nonlinear stochastic-dynamical model. *Phys Rev E* 2012, 85. doi: 10.1103/PhysRevE.85.031134.
82. Franzke C. Predictions of critical transitions with non-stationary reduced order models. *Phys D* 2013, 262:35–47.
83. Sura P. A general perspective of extreme events in weather and climate. *Atmos Res* 2011, 101:1–21.
84. Sura P. Stochastic models of climate extremes: theory and observations. In: AghaKouchak A, Easterling D, Hsu K, Schubert S, Sorooshian S, eds. *Extremes in a Changing World: Detection, Analysis and Uncertainty*. Berlin: Springer; 2013, 432.
85. Heinz S. *Statistical Mechanics of Turbulent Flows*. Berlin: Springer; 2003, 215.
86. Kraichnan RH. The structure of isotropic turbulence at very high reynolds number. *J Fluid Mech* 1959, 5:497.
87. Frederiksen JS. Subgrid-scale parameterizations of eddy-topographic force, eddy viscosity, and stochastic backscatter for flow over topography. *J Atmos Sci* 1999, 56:1481.

88. Herring JR, Kraichnan RH. *Statistical Models of Turbulence*. Berlin: Springer; 1972.
89. Leith CE. Atmospheric predictability and two-dimensional turbulence. *J Atmos Sci* 1971, 28:145.
90. Leith CE, Kraichnan RH. Predictability of turbulent flows. *J Atmos Sci* 1972, 29:1041–1058.
91. Lorenz EN. A study of the predictability of a 28-variable atmospheric model. *Tellus* 1965, 17:321–333.
92. Frederiksen JS, O’Kane TJ. Inhomogeneous closure and statistical mechanics for Rossby wave turbulence over topography. *J Fluid Mech* 2005, 539:137.
93. O’Kane TJ, Frederiksen JS. A comparison of statistical dynamical and ensemble prediction methods during blocking. *J Atmos Sci* 2008b, 65:426.
94. Epstein ES. Stochastic dynamic prediction. *Tellus* 1969, 21:739–759.
95. Pitcher EJ. Application of stochastic dynamic prediction to real data. *J Atmos Sci* 1977, 34:3–21.
96. Hunte MWDA, Johnson C. Conditional mode elimination and the subgrid-modelling problem for isotropic turbulence. *Phys Fluids* 2001a, 13:2030.
97. Frederiksen JS, Davies AG. Eddy viscosity and stochastic backscatter parameterizations on the sphere for atmospheric circulation models. *J Atmos Sci* 1997, 54:2475.
98. Kraichnan RH. Eddy viscosity in two and three dimensions. *J Atmos Sci* 1976, 33:1521.
99. Leslie DC, Quarini GL. The application of turbulence theory to the formulation of subgrid modelling procedures. *J Fluid Mech* 1979, 91:65.
100. O’Kane TJ, Frederiksen JS. Statistical dynamical subgrid-scale parameterizations for geophysical flows. *Phys Scr* 2008a, T132:014033.
101. Rose HA. Eddy diffusivity, eddy noise and subgrid-scale modelling. *J Fluid Mech* 1977, 81:719.
102. Kitsios V, Frederiksen JS, Zidikheri MJ. Subgrid model with scaling laws for atmospheric flows. *J Atmos Sci* 2012, 69:1427.
103. Frederiksen JS, Davies AG. Dynamics and spectra of cumulant update closures for two-dimensional turbulence. *Geophys Astrophys Fluid Dyn* 2000, 92:197.
104. O’Kane TJ, Frederiksen JS. The QDIA and regularized QDIA closures for inhomogeneous turbulence over topography. *J Fluid Mech* 2004, 504:133.
105. Herring JR. Self-consistent-field approach to turbulence theory. *Phys Fluids* 1965, 8:2219.
106. McComb WD. A local energy-transfer theory of isotropic turbulence. *J Phys A* 1974, 7:632.
107. Carnevale GF, Frederiksen JS. Viscosity renormalization based on direct-interaction closure. *J Fluid Mech* 1983, 131:289.
108. Dekker U, Haake F. Fluctuation-dissipation theorems for classical processes. *Phys Rev A* 1957, 6:2043.
109. Frederiksen JS, Davies AG, Bell RC. Closure equations with non-Gaussian restarts for truncated two-dimensional turbulence. *Phys Fluids* 1994, 6:3153.
110. Ogura Y. A consequence of the zero fourth cumulant approximation in the decay of isotropic turbulence. *J Fluid Mech* 1963, 16:33.
111. Orszag SA. Analytical theories of turbulence. *J Fluid Mech* 1970, 41:363.
112. Leith CE. Theoretical skill of Monte Carlo forecasts. *Mon Weather Rev* 1974, 102:409.
113. Métais O, Lesieur M. Statistical predictability of decaying turbulence. *J Atmos Sci* 1986, 43:857.
114. O’Kane TJ, Frederiksen JS. Comparison of statistical dynamical, square root and ensemble Kalman filters. *Entropy* 2008c, 10:684–721. doi: 10.3390/e10040684.
115. O’Kane TJ, Frederiksen JS. The application of statistical dynamical turbulence closures to data assimilation. *Phys Scr* 2010, T142:014042.
116. Rose HA. An efficient non-Markovian theory of non-equilibrium dynamics. *Phys D* 1985, 14:216.
117. Memin E. Fluid flow dynamics under location uncertainty. *Geophys Astrophys Fluid Dyn* 2014, 108:119–146.
118. Smagorinsky J. General circulation experiments with the primitive equations, I: the basic experiment. *Mon Weather Rev* 1963, 91:99.
119. Holloway G. Representing topographic stress for large-scale ocean models. *J Phys Oceanogr* 1992, 22:1033.
120. Zidikheri MJ, Frederiksen JS. Stochastic subgrid parameterizations for simulations of atmospheric baroclinic flows. *J Atmos Sci* 2009, 66:2844.
121. Palmer TN. A nonlinear dynamical perspective on model error: a proposal for non-local stochastic-dynamic parameterization in weather and climate prediction. *Q J Roy Meteorol Soc* 2001, 127:279–304.
122. Penland C. Noise out of chaos and why it won’t go away. *Bull Am Meteorol Soc* 2003, 84:921–925.
123. Sardeshmukh P, Penland C, Newman M. Rossby waves in a stochastically fluctuating medium. In: Imkeller P, von Storch J-S, eds. *Stochastic Climate Models*. Basel: Birkhäuser; 2001, 384–396.
124. Berner J, Jung T, Palmer TN. Systematic model error: the impact of increased horizontal resolution versus improved stochastic and deterministic parameterizations. *J Climate* 2012, 25:4946–4962.
125. Cohen BG, Craig GC. Fluctuations in an equilibrium convective ensemble, part II: numerical experiments. *J Atmos Sci* 2005, 63:2005–2015.
126. Plant RS, Craig GC. A stochastic parameterization for deep convection based on equilibrium statistics. *J Atmos Sci* 2008, 65:87–105.

127. Hagedorn R, Doblas-Reyes FJ, Palmer TN. The rationale behind the success of multi-model ensembles in seasonal forecasting—i. basic concept. *Tellus* 2005, 57A:219–233.
128. Houtekamer PL, Lefaiivre L, Derome J, Ritchie H, Mitchell HL. A system simulation approach to ensemble prediction. *Mon Weather Rev* 1996, 124:1225–1242.
129. Krishnamurti TN, Kishtawal CM, Zhang Z, LaRow T, Bachiochi D, Williford E, Gadgil S, Surendran S. Multimodel ensemble forecasts for weather and seasonal climate. *J Climate* 2000, 13:4196–4216.
130. Stainforth D, Aina T, Christensen C, Collins M, Faull N, Frame DJ, Kettleborough JA, Knight S, Martin A, Murphy JM, et al. Uncertainty in predictions of the climate response to rising levels of greenhouse gases. *Nature* 2005, 433:403–406.
131. Berner J, Ha S-Y, Hacker JP, Fournier A, Snyder C. Model uncertainty in a mesoscale ensemble prediction system: stochastic versus multi-physics representations. *Mon Weather Rev* 2011, 139:1972–1995.
132. Eckel FA, Mass CF. Aspects of effective mesoscale, short-range ensemble forecasting. *Wea Forecasting* 2005, 20:328–350.
133. Murphy JM, Sexton DM, Barnett DN, Jones GS, Webb MJ, Collins M, Stainforth DA. Quantification of modelling uncertainties in a large ensemble of climate change simulations. *Nature* 2004, 430:768–772.
134. Stensrud DJ, Bao J-W, Warner TT. Using initial condition and model physics perturbations in short-range ensemble simulations of mesoscale convective systems. *Mon Weather Rev* 2000, 128:2077–2107.
135. Palmer T, Andersen U, Cantelaube P, Davey M, Deque M, Doblas-Reyes F, Feddersen H, Graham R, Gualdi S, Gueremy J-F, et al. Development of a European multi-model ensemble system for seasonal to inter-annual prediction (DEMETER). *Bull Am Meteorol Soc* 2004, 85:853–872.
136. Stocker ETF, Qin D, Plattner G-K, Tignor M, Allen S, Boschung J, Nauels A, Xia Y, Bex V, Midgley P. *Climate Change 2013: The Physical Science Basis*. Cambridge; New York: Cambridge University Press; 2013.
137. Bowler NE, Arribas A, Beare SE, Mylne KR, Shutts GJ. The local ETKF and SKEB: upgrades to the MOGREPS short-range ensemble prediction system. *Q J Roy Meteorol Soc* 2009, 135:767–776.
138. Palmer, TN, Buizza R, Doblas-Reyes F, Jung T, Leutbecher M, Shutts G, Steinheimer M, Weisheimer A. Stochastic parametrization and model uncertainty. ECMWF Technical Memorandum No. 598, 2009. Available at: <http://www.ecmwf.int/publications/>.
139. Berner J, Doblas-Reyes FJ, Palmer TN, Shutts G, Weisheimer A. Impact of a quasi-stochastic cellular automaton backscatter scheme on the systematic error and seasonal prediction skill of a global climate model. *Phil Trans R Soc A* 2008, 366:2559–2577.
140. Doblas-Reyes F, Weisheimer A, Déqué M, Keenlyside N, McVean M, Murphy JM, Rogel P, Smith D, Palmer TN. Addressing model uncertainty in seasonal and annual dynamical seasonal forecasts. *Q J Roy Meteorol Soc* 2009, 135:1538–1559.
141. Weisheimer A, Palmer T, Doblas-Reyes F. Assessment of representations of model uncertainty in monthly and seasonal forecast ensembles. *Geophys Res Lett* 2011, 38:L16703. doi: 10.1029/2011GL048123.
142. Epstein ES, Pitcher EJ. Stochastic analysis of meteorological fields. *J Atmos Sci* 1972, 29:244–257.
143. Lorenz EN. Deterministic nonperiodic flow. *J Atmos Sci* 1963, 20:130–141.
144. Molteni F, Palmer TN. Predictability and finite-time instability of the Northern winter circulation. *Q J Roy Meteorol Soc* 1993, 119:269–298.
145. Toth Z, Kalnay E. Ensemble forecasting at NMC: the generation of perturbations. *Bull Am Meteorol Soc* 1993, 74:2317–2330.
146. Buizza R, Houtekamer PL, Toth Z, Pellerini G, Wei M, Zhu Y. A comparison of the ECMWF, MSC and NCEP global ensemble prediction systems. *Mon Weather Rev* 2005, 133:1076–1097.
147. Palmer TN. A nonlinear dynamical perspective on climate prediction. *J Climate* 1999, 12:575–591.
148. Smith LA. *Disentangling Uncertainty and Error: On the Predictability of Nonlinear Systems*. Berlin: Springer; 2001.
149. Buizza R, Miller M, Palmer TN. Stochastic representation of model uncertainties in the ECMWF ensemble prediction system. *Q J Roy Meteorol Soc* 1999, 125:2887–2908.
150. Berner, J, Smith KR, Ha S-Y, Hacker JP, and Snyder C. Increasing the skill of probabilistic forecasts: model-error representations versus calibration and debiasing. *Mon Weather Rev*. Submitted for publication.
151. Reynolds CA, McLay JG, Goerss JS, Serra EA, Hodyss D, Sampson CR. Impact of resolution and design on the U.S. Navy global ensemble performance in the tropics. *Mon Weather Rev* 2011, 139:2145–2155.
152. Teixeira J, Reynolds CA. Stochastic nature of physical parameterization in ensemble prediction: a stochastic convection approach. *Mon Weather Rev* 2008, 136:483–496.
153. Mason P, Thomson D. Stochastic backscatter in large-eddy simulations of boundary layers. *J Fluid Mech* 1992, 242:51–78.
154. Shutts GJ, Palmer TN. The use of high resolution numerical simulations of tropical circulation to calibrate stochastic physics schemes. In: *ECMWF/CLIVAR Workshop on Simulation and Prediction of Intra-seasonal Variability with the*

- Emphasis on the MJO*, ECMWF, 2003, 83–102. Available at: <http://www.ecmwf.int/publications>.
155. Bowler NE, Arribas A, Mylne KR, Robertson KB, Beare SE. The MOGREPS short-range ensemble prediction system. *Q J Roy Meteorol Soc* 2008, 134:703–722.
 156. Charron M, Pellerin G, Spacek L, Houtekamer PL, Gagnon N, Mitchell HL, Michelin L. Toward random sampling of model error in the Canadian ensemble prediction system. *Mon Weather Rev* 2010, 138:1877–1901.
 157. Sanchez, C, Williams KD, Shutts G, Collins M, Impact of a stochastic kinetic energy backscatter scheme across time-scales and resolutions. *Q J Roy Meteorol Soc* 2014, 140:577–599. doi:10.1002/qj.2328.
 158. Bouttier F, Vié B, Nuissier O, Raynaud L. Impact of stochastic physics in a convection-permitting ensemble. *Mon Weather Rev* 2012, 140:3706–3721.
 159. Romine GS, Schwartz CS, Berner J, Smith KR, Snyder C, Anderson JL, and Weisman ML: Representing forecast error in a convection-permitting ensemble system. *Mon Weather Rev*. Submitted for publication.
 160. Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM. *Climate Change 2013: The Physical Science Basis*. Cambridge: Cambridge University Press; 2013, 1535.
 161. Lin JWB, Neelin JD. Influence of a stochastic moist convective parameterization on tropical climate variability. *Geophys Res Lett* 2000, 27:3691–3694.
 162. Frenkel Y, Majda AJ, Khouider B. Using the stochastic multicloud model to improve tropical convective parameterizations: a paradigm example. *J Atmos Sci* 2011, 69:1080–1105.
 163. Khouider B, Biello J, Majda AJ. A stochastic multicloud model for tropical convection. *Commun Math Sci* 2010, 8:187–216.
 164. Majda AJ, Stechmann S. Stochastic models for convective momentum transport. *Proc Natl Acad Sci USA* 2008, 105:17614–17619.
 165. Jung T, Palmer TN, Shutts GJ. Influence of a stochastic parameterization on the frequency of occurrence of north pacific weather regimes in the ECMWF model. *Geophys Res Lett* 2005, 32:L23811.
 166. Palmer TN, Weisheimer A. Diagnosing the causes of bias in climate models—why is it so hard? *Geophys Astrophys Fluid Dyn* 2011, 105:351–365.
 167. Williams PD. Climatic impacts of stochastic fluctuations in air-sea fluxes. *Geophys Res Lett* 2012, 39:L10705. doi: 10.1029/2012GL051813.
 168. Sapsis T, Majda AJ. A statistically accurate modified quasilinear Gaussian closure for uncertainty quantification in turbulent dynamical systems. *Phys D* 2013a, 252:34–45.
 169. Sapsis T, Majda AJ. Statistically accurate low-order models for uncertainty quantification in turbulent dynamical systems. *Proc Natl Acad Sci USA* 2013b, 110:13705–13710.
 170. Solomon SE, Qin D, Manning M, Chen Z, Marquis M, Averyt K, Tignor M, Miller H. *Climate Change 2007: The Physical Science Basis*. New York: Cambridge University Press; 2007, 996.
 171. Palmer T. More reliable forecasts with less precise computations: a fast-track route to cloud-resolved weather and climate simulators? *Phil Trans R Soc A* 2014, 372:20130391.
 172. Vanden-Eijnden E. Numerical techniques for multi-scale dynamical systems with stochastic effects. *Commun Math Sci* 2003, 1:385–391.
 173. Gottwald G, Harlim J. The role of additive and multiplicative noise in filtering complex dynamical systems. *Proc R Soc A* 2013, 469:20130096.
 174. Ha, S-Y, J Berner, and C Snyder: Model error representation in mesoscale wrf-dart cycling. *Mon Weather Rev*. Submitted for publication.
 175. Isaksen, L, Fisher M, Berner J. Use of analysis ensembles in estimating flow-dependent background error variances. In: *Proceedings of ECMWF Workshop on Flow Dependent Background Errors* ECMWF, Reading, UK, 11–13 June, 2007, 65–86.
 176. Miller RN, Carter EF, Blue ST. Data assimilation into nonlinear stochastic models. *Tellus* 1999, 51A:167–194.
 177. Mitchell L, Gottwald GA. Data assimilation in slow-fast systems using homogenized climate models. *J Atmos Sci* 2012, 69:1359–1377.

FURTHER READING

- Franzke C, Majda AJ, Branstator G. The origin of nonlinear signatures of planetary wave dynamics: mean phase space tendencies and contributions from non-Gaussianity. *J Atmos Sci* 2007, 64:3987–4003. doi: 10.1175/2006JAS2221.1.
- Kraichnan RH. Test-field model for inhomogeneous turbulence. *J Fluid Mech* 1972, 56:287.
- Shutts G, Allen T, Berner J. Stochastic parametrization of multiscale processes using a dual-grid approach. *Phil Trans R Soc A* 2008, 366:2623–2639.